arXiv:1304.2910v1 [quant-ph] 10 Apr 2013

Reliable quantum replication at the Heisenberg limit

Giulio Chiribella*
Center for Quantum Information, Institute for Interdisciplinary Information Sciences, Tsinghua University, Beijing, 100084, China

Yuxiang Yang†
Department of Physics, Tsinghua University Beijing, 100084, China

Andrew Chi-Chih Yao‡
Institute for Interdisciplinary Information Sciences, Tsinghua University, Beijing, 100084, China§

No process in nature can perfectly clone an arbitrary quantum state. But is it possible to engineer processes that replicate quantum information with vanishingly small error? Here we demonstrate the possibility of probabilistic super-replication phenomena where N equally prepared quantum systems are transformed into a much larger number M nearly perfect replicas, with an error that rapidly vanishes whenever M is small compared to N^2. The quadratic replication rate is the ultimate limit imposed by quantum mechanics and is fundamentally linked with the Heisenberg limit of quantum metrology.

No physical process can copy arbitrary quantum states on demand [1, 2]: if such a process existed, we could build a device that distinguishes quantum states with arbitrary precision, violating the uncertainty principle and enabling faster-than-light communication [3]. Probabilistic processes like stimulated emission, however, seem to evade this restriction. An excited atom interacting with a polarized photon is expected to produce sometimes a second photon with the same polarization, effectively delivering a perfect clone. True that some other times the atom will spontaneously emit a photon of random polarization, but still, when stimulated emission occurs, a prefect clone has been produced. If we had on our side a quantum version of Maxwell demon, who separates the photons produced by stimulated emission from those produced by spontaneous emission, we would be able to produce any desired number of clones with a non-zero probability. Unfortunately, our imaginary helper is not allowed by the laws of quantum mechanics: even with small probability, it would allow us to violate the no-signalling principle [4]. However, nothing forbids that other probabilistic processes, akin to spontaneous emission, could boost our ability to replicate quantum information beyond any previously conceived limit. This possibility raises new fundamental questions: Is it possible, for example, to engineer a process that duplicates a beam of N equally prepared particles, producing a beam of $2N$ almost perfect clones? What is the ultimate limit to the number of replicas that can be produced without significant errors by a physical process?

Here we answer both questions: For a beam of N quantum particles in an arbitrary state generated by Hamiltonian evolution, we devise a filtering mechanism that allows one not only to duplicate, but also to produce an overwhelming number M of nearly perfect clones, with a small error that vanishes rapidly whenever M is small compared to N^2. We name this new phenomenon *super-replication* and show that it is intrinsically probabilistic, proving that deterministic processes can only produce a negligible number of nearly perfect replicas. For example, for 100 linearly polarized photons, super-replication allows to produce 1000 replicas with fidelity 99.9%, whereas the best deterministic process can only achieve without filter the fidelity 57%. In addition, we show that no physical process, deterministic or not, can produce more than $O(N^2)$ replicas: any attempt to replicate quantum information beyond this limit is doomed to produce a joint output that has vanishing fidelity with the desired state. To explain the roots of this fundamental limitation, we establish a deep link between quantum cloning and the precision limits of quantum metrology [5–10], showing that the Heisenberg Limit sets the ultimate bound to the replication rate of probabilistic processes, while the Standard Quantum Limit sets the corresponding bound for deterministic processes.

I. RESULTS

Quantum metrology limits to information replication. The *Heisenberg Limit (HL)* is the cornerstone of quantum metrology [5–10]. It states that whenever one tries to estimate the value of a parameter characterizing a physical process, the variance of the estimate will decrease at most as $1/N^2$, where N is the number of times that the process is probed. The scaling $1/N^2$ can be achieved, provided that the probes are prepared in a suitable entangled state. However, if the probes are not entangled, then the variance decreases only as $1/N$—

---

[*] gchiribella@mail.tsinghua.edu.cn
[†] yangyx09@mails.tsinghua.edu.cn
[‡] andrewcyao@mail.tsinghua.edu.cn
[§] http://iiis.tsinghua.edu.cn

known as the *Standard Quantum Limit (SQL)*, which is the scaling of a classical statistics over repeated experiments.

At first sight, the Heisenberg Limit seems to have nothing to do with quantum cloning: the former limits the quality of estimation using entangled states, the latter aims at producing copies from a product state in input. However, we will now show that the HL sets the ultimate upper bound on the performance of cloning. We consider the setting where N identical systems are given, and each system is prepared in the same state $|\psi_\theta\rangle = e^{i\theta H}|\psi\rangle$, for some Hamiltonian H and for some unknown parameter $\theta \in \mathbb{R}$. For example, θ could be the polarization angle of a linearly polarized photon, or the time parameter in a dynamics governed by the Schrödinger's equation. To link quantum cloning and quantum metrology, we adopt an asymptotic approach inspired by information theory: we consider cloning processes that transform N input copies into $M = cN^\alpha$ approximate copies, where $c > 0$ and $\alpha > 1$ are suitable constants. We call the constant α the *replication rate* and say that a process achieves *reliable replication* if the error vanishes in the limit of large N.

Now the question is: how many copies of the state $|\psi_\theta\rangle$ can be produced by a reliable replication process? The answer, argued for in the Methods section, is given by the following:

1. **Heisenberg Limit for Quantum Replication:** *No physical process can reliably replicate the quantum states $\{e^{i\theta H}|\psi\rangle, \theta \in \mathbb{R}\}$ at a rate larger than* 2.

The Heisenberg Limit leaves lots of room for replicating quantum information: for every rate $1 < \alpha < 2$ one has the chance not only to duplicate the input copies, but also to produce an overwhelming number of replicas. In the next paragraph, we will exhibit explicit schemes that achieve replication at every rate allowed by the HL. These schemes are necessarily probabilistic, as a consequence of the Standard Quantum Limit: in the Methods section we show that the SQL implies that every deterministic process is bound to produce a number of replicas that is negligible with respect to N:

2 **Standard Quantum Limit for Quantum Replication:** *No deterministic process can reliably replicate the states $\{e^{i\theta H}|\psi\rangle, \theta \in \mathbb{R}\}$ at a rate larger than* 1.

In other words, without a probabilistic filter, one can only embezzle from nature a negligible number of extra-copies.

The difference between probabilistic and deterministic processes does not consist only in the number of replicas that can be produced reliably, but also in the *speed* at which the replication error vanishes: as proven in the Methods section, the error cannot vanish faster than (the inverse of) a polynomial of degree smaller than 4 in the deterministic case, whereas in the probabilistic case the error can vanish faster than every polynomial.

Probabilistic super-replication. Here we show that the states $|\psi_\theta\rangle = e^{i\theta H}|\psi\rangle$ can be reliably replicated at any rate allowed by the HL. When the rate is larger than (or equal to) 1, we call the process *super-replication* to emphasize the fact that it beats the SQL. In addition, we pin down the probabilistic nature of super-replication, showing a tight relation between the replication rate and the probability that replication takes place successfully.

The key idea to achieve super-replication is to devise a quantum filter that modulates the Fourier amplitudes of the wavefunction, in a way that enhances the cloning performances to the maximum rate allowed by quantum mechanics. To move to the Fourier picture, we express the input states as $|\psi_\theta\rangle = \sum_{E\in\mathsf{Spec}(H)} \sqrt{p_E} e^{i\theta E}|E\rangle$, where $\mathsf{Spec}(H)$ denotes the spectrum of the Hamiltonian, $|E\rangle$ is an eigenvector of H with eigenvalue E, and p_E is the probability that a measurement of the energy gives outcome E. Without loss of generality, we assume that the average energy is zero, a condition that can always be enforced by shifting all the eigenvalues by a suitable constant. When N identical copies are given, the joint state can be expressed as

$$|\psi_\theta\rangle^{\otimes N} = \sum_{E\in\mathsf{Spec}(H^{(N)})} e^{i\theta E}\sqrt{p_{N,E}}|N,E\rangle, \qquad (1)$$

where $H^{(N)} = \sum_{i=1}^N H_i$ is the total Hamiltonian (H_i denoting the one-body Hamiltonian H for the i-th system), $|N,E\rangle$ is an eigenvector of $H^{(N)}$ for the eigenvalue E, and $p_{N,E}$ is the probability that a measurement of the total energy gives outcome E.

The ideal goal of cloning is to produce M perfect copies in the state $|\psi_\theta\rangle^{\otimes M}$. The filter is described by two measurement operators M_{yes} and M_{no}, acting on the space of N copies. When the N input systems emerge from the filter they are in the state $M_{yes}|\psi_\theta\rangle^{\otimes N}$, up to a normalization factor. Now, by suitably choosing the operator M_{yes}, we can modulate the Fourier amplitudes of the input state in any way we like. One natural choice would be to replace N with M in the probability distribution $p_{N,E}$. Since, however, the spectrum of $H^{(N)}$ may be not be contained in the spectrum of $H^{(M)}$, we may need also to shift the values of the energy by a small amount E_0, which tends to zero for large M (see the Supplemental Material for details). With this choice, the state after the filter is projected to the entangled state $|\Psi_\theta^N\rangle \propto \sum_{E\in\mathsf{Spec}(H^{\otimes N})} e^{i\theta E}\sqrt{p_{M,E+E_0}}|N,E\rangle$.

The state $|\Psi_\theta^N\rangle$ will now act as a *quantum program*, containing the instructions that will be used by a deterministic quantum device to generate M approximate clones. For this purpose, we consider a quantum device that coherently transforms each basis state $|N,E\rangle$ into the corresponding state $|M,E+E_0\rangle$, thus producing the state $|\Phi_\theta^M\rangle \propto \sum_{E\in\mathsf{Spec}(H^{(N)})} e^{i\theta E}\sqrt{p_{M,E+E_0}}|M,E+E_0\rangle$. It is not hard to see that the state $|\Phi_\theta^M\rangle$ has a very high overlap with the desired state $|\psi_\theta\rangle^{\otimes M}$ for every replication rate allowed by the Heisenberg limit. For large N, using the central limit theorem one can show that the fidelity satisfies the following lower bound

$$F_{N\to M} \geq 1 - \sqrt{\frac{M\langle H^2\rangle}{\pi\eta N^2}}\exp\left(-\frac{\eta N^2}{M\langle H^2\rangle}\right), \qquad (2)$$

where η is a suitable constant (see Supplemental Material). The bound shows that the fidelity rapidly approaches 1 whenever M is of order N^α with α smaller than 2, that is, whenever the replication rate satisfies the Heisenberg Limit. A strong converse can be proven too: every physical process replicating quantum states at a rate higher than 2 must have vanishing fidelity in the limit of large N.

Thanks to the use of our filter, we have been able to embezzle from nature a large number of replicas. The improvement is striking if one compares it with the performances of standard cloning processes. Let us illustrate this fact for the replication of linear polarization states $|\psi_\theta\rangle = \cos\theta|V\rangle + \sin\theta|H\rangle$. In this case, the best deterministic process is the phase-covariant cloner of Ref. [11], and its fidelity is $F^{det}_{N\to M} \approx \frac{2\sqrt{MN}}{M+N}$ in the asymptotic limit of large N and M. In agreement with our Standard Quantum Limit, the fidelity vanishes whenever M is of order $N^{1+\epsilon}$, $\epsilon > 0$: As we proved in the general case, deterministic processes can only produce a number of replicas that is a negligible fraction of the number of input photons. For example, starting 100 from input photons, a deterministic process can produce only a few extra-copies, while using our filter one can produce up to thousands nearly perfect replicas. For $N = 100$ and $M = 1000$, cloning with filter achieves fidelity $F_{prob} = 0.9986$, while without filter the fidelity is $F_{det} = 0.5739$. The different features of replication processes at the Heisenberg limit and at the standard quantum limit are illustrated in Fig. 1. A plot comparing the performances of replication with and without the filter in the case $N = 20$ is presented in Fig. 2. What makes the improvement even more dramatic is that the quality of the replicas is measured by the *global fidelity* between the output state and the desired joint state of M perfect copies, which is much smaller than the overlap that each single copy has with the state $|\psi_\theta\rangle$.

Maximizing the probability of successful replication. The dramatic enhancement leading to super-replication comes at a price: the probability that the input systems pass the filter has to decay with N. For our filter, the probability decreases exponentially fast as $p_{yes}[N \to M] \leq e^{-kN}$ for a suitable constant $k > 0$. One can ask whether there are ways to achieve super-replication with a larger success probability. It turns out that the answer is *yes*, and that larger success probabilities can be obtained through a minimal modification of our filter. In particular, one can generate $M = cN$ replicas with a probability $p_{yes}[N \to M]$ going to zero as $1/N^\delta$ for every desired $\delta > 0$. The price to pay, however,



FIG. 1. **More and better: the advantages of probabilistic super-replication** (a) Every deterministic process is constrained by the Standard Quantum Limit: the number of clones that can be produced reliably from N input copies is of order N and the fidelity of the clones cannot approach 1 faster than the inverse of a polynomial of degree 4. (b) Super-replication at the Heisenberg limit: the cloning performances can be dramatically increased by a probabilistic filter, depicted here as a "quantum Maxwell demon" that separates two branches of the wavefunction. In the successful branch, any number of clones of order $N^{2-\epsilon}$ can be produced with fidelity approaching 1 faster than any polynomial. The probabilistic filter allows one to generate more clones and with a better quality.



FIG. 2. **Replication of linearly polarized photons: with and without filter.** The fidelity of the best cloning processes for linearly polarized photons $|\psi_\theta\rangle = \cos\theta|V\rangle + \sin\theta|H\rangle$ is plotted here for $N = 20$ input copies, with the number M of output clones ranging from 20 to 400. The red (green) line refers to optimal cloning with (without) filter.

is that the fidelity will go to 1 only as $1/(N^{\delta}\ln N)$. More generally, in the Supplemental Material we show that every replication process with non-vanishing probability must necessarily have vanishing fidelity. This provides a strong converse of the Standard Quantum Limit: every deterministic process producing replicas at a rate higher than the SQL must have vanishing fidelity.

For super-replication processes with rate $\alpha = 1+\epsilon$, $\epsilon > 0$, we show how the success probability can be increased to $p_{yes}[N \to M] \geq e^{-N^{\delta}}$ for every desired exponent $\delta > \epsilon$. However, no further improvement is possible below the critical value $\delta_c = \epsilon$: any process with success probability scaling as $p_{yes}[N \to M] = ae^{-bN^{\delta}}$, $\delta < \epsilon$ and for $a, b > 0$ must have vanishing fidelity in the asymptotic limit.

Delayed choice between deterministic and probabilistic strategies. If information is money, an easy way to picture the difference between deterministic and probabilistic replication is as two different ways to make investments. Deterministic processes are like depositing money in a bank with low interest (one gets a negligible amount of extra replicas), whereas probabilistic processes are like buying tickets of a lottery with large jackpot (the prize is big, but the chance to win is small). Depending on the situation, one or the other approach may be more convenient. Interestingly, a bonus of our results is that one can always delay the choice whether or not to use the filter. Indeed, one can first implement the deterministic transformation $|N,E\rangle \mapsto |M, E+E_0\rangle$ on the energy eigenstates and postpone the modulation of the probability distribution to the very end. At this stage, if the modulation is omitted, one has a nearly optimal deterministic process for replication at the SQL. Instead, if the modulation is performed, one can probabilistically enhance the replication performance to the Heisenberg-limit scaling.

The many-worlds fairness of quantum mechanics. Super-replication can be achieved not only for one-parameter families of states of the form $|\psi_\theta\rangle = e^{i\theta H}|\psi\rangle$, but also for different manifolds of states, including the manifold of all maximally entangled states of two identical systems. However, super-replication is not a generic feature. For example, if one tries to copy an *arbitrary*—as opposed to linear—polarization state, then no filter is going to help: the performances with filter are equal to the performances without filter. What is the origin of such a difference between linear and arbitrary polarization? As we show in the Supplemental Material, the explanation comes from the geometry of quantum states: states on a circle can be replicated at the Heisenberg limit, but states on the sphere cannot. We also show that such a limitation holds for quantum systems of arbitrary dimension: *no probabilistic filter can make an arbitrary quantum state more copiable.* This property is quite compelling when considered from the angle of the many-worlds interpretation of quantum mechanics, because it states that no-branch of the wavefunction of the universe offers an advantage over the others when it comes to replicating the information contained in a completely unknown state. Regarding different branches as "different worlds", we can formulate this as a fundamental principle, which we name *many-worlds fairness*, stating that the capability to replicate arbitrary information is the same in all possible worlds. Many world fairness rules out quantum mechanics on *real* Hilbert spaces [12], an alternative physical theory where photons can have only linear polarizations. Thanks to this observation, one can provide a new answer to the old question"Why are physical systems described by *complex*,—instead of *real* Hilbert spaces?". Traditionally, the standard answer has been to invoke *local tomography*, namely the property that one can completely identify a mixed state from the statistics of local measurement on the components. However, one may consider it as an *ad hoc* requirement [13], and, in fact, there are even reasons to prefer real quantum mechanics to its complex version, as it was recently pointed out by Wotters [14]. Balancing this fact, many world fairness offers a new reason (other that the usual local tomography) to prefer complex quantum mechanics to its real relative.

II. DISCUSSION

Towards experimental implementation. Super-replication at the Heisenberg limit has been introduced here from a theoretical point of view. But is it possible to implement it experimentally? Luckily, very recently there have been important experimental breakthroughs on the closely related topic of probabilistic amplification of coherent states of light [15–19]. Although the translation to our case is not immediate, we suggest that super-replication of linearly polarized photons could be achieved through a suitable sequence of amplitude damping channels, which for the polarization play the role of the photon subtraction for coherent states. An alternative approach would be to first encode the states of the input copies into coherent states via matter-light teleportation [20], amplify the latter, and then teleport back. This would provide a new application to the existing experimental schemes of coherent state amplification, making them the building block for the replication of quantum information at the Heisenberg limit.

Relation with previous works on probabilistic cloning. The idea that in some situations even non-orthogonal states can be copied perfectly by a probabilistic device was first introduced by Duan and Guo [21], who showed that a set of pure quantum states can be copied perfectly if and only if they are linearly independent. This means that for a single photon one can perfectly copy two polarization states, while for N photons one can copy at most $N+1$ states. Subsequent works [22, 23] showed that probabilistic devices can improve the performances of approximate cloning, leading to nearly ideal performances in the case of coherent states with equal amplitude. The nearly perfect cloning of coherent states may lead one to believe that there is no limit to the amount of clones that can be produced probabilistically. Contrarily to this intuition, we have proven here that the Heisenberg limit sets the ultimate bound $M = O(N^2)$ to the number of clones that can be produced reliably. Moreover, we have shown that for arbitrary one-parameter families of states, any replication rate allowed by the Heisenberg Limit can be achieved, and that the process that achieves it must be probabilistic, because deterministic processes are constrained by the Standard Quantum Limit $M = O(N)$.

Relation with probabilistic quantum metrology. Recently it has been observed that the precision in the estimation of a parameter encoded in a set of quantum states can be improved using probabilistic schemes [24, 25] where, depending on the outcome of the filter, one can decide to *abstain* from estimating the parameter. The mechanism of abstention can sometimes boost the precision of estimation from the standard quantum limit $1/N$ to the Heisenberg limit $1/N^2$. The connection between these results and the super-replication phenomenon is made clear by our approach to the Heisenberg limit: the fact that a probabilistic filter can improve estimation until the Heisenberg limit implies that N^2, rather than N, is the upper bound for the replication rate of the states $|\psi_\theta\rangle = e^{i\theta H}|\psi\rangle$. However, there is an important catch: *the filter that achieves replication at the Heisenberg limit is not the same filter that achieves estimation at the Heisenberg limit.* In fact, if we were to use the same filter needed in Refs. [24, 25], we would not be able to reduce the error down to zero.

Conclusions. We discovered a previously unexplored connection between quantum cloning and quantum metrology, showing the Standard Quantum Limit and the Heisenberg Limit for estimation can be translated into limits for the reliable replication of quantum information. The possibility of producing a large number of almost perfect replicas, opened by the Heisenberg Limit, has been demonstrated for arbitrary one-parameter families of states generated by a Hamiltonian H, showing that, with the help of a probabilistic filter, one can embezzle from nature a large number number of extra clones, as long as the number is small compared to N^2, the square of the number of input copies. We then showed that the phenomenon of super-replication is not generic: In particular, there is no way to improve the performances of cloning of an arbitrary polarization state. The fact that the geometry of quantum states forbids any advantage in the replication of an arbitrary state based on the probabilistic branching of the wavefunction is a fundamental property of quantum mechanics, that we call *many-world fairness.* It is our hope that our work will stimulate the debate on the foundations of quantum mechanics and trigger new advancements on the experimental realization of probabilistic quantum information-processing devices.

III. METHODS

Decomposition of quantum instruments. To describe the most general probabilistic processes allowed by quantum mechanics, we use the framework of quantum instruments. A quantum instrument with input (output) Hilbert space $\mathscr{H}_{in}$ ($\mathscr{H}_{out}$) is a collection of completely positive, trace non-increasing linear maps $\{\mathcal{P}_i\}_{i\in\mathsf{X}}$, each map transforming density matrices on $\mathscr{H}_{in}$ into subnormalized density matrices on $\mathscr{H}_{out}$. If the input system is prepared in the density matrix ρ, the probability that the i-th process takes place is $p(i|\rho) = \operatorname{Tr}[\mathcal{P}_i(\rho)]$. Conditionally to the occurrence of outcome i, the state of the output system is $\rho'_i = \mathcal{P}_i(\rho)/\operatorname{Tr}[\mathcal{P}_i(\rho)]$. We note an elementary fact about quantum instruments: *any quantum instrument can be decomposed into a pure measurement on the input system followed by quantum channel depending on the outcome.* By "pure measurement on the input system", we mean a quantum instrument $\{\mathcal{M}_i\}_{i\in\mathsf{X}}$ of the special form $\mathcal{M}_i(\rho) = M_i\rho M_i^\dagger$ for some measurement operator M_i on $\mathscr{H}_{in}$. By "quantum channel", we mean a completely positive trace-preserving map $\mathcal{C}_i$, transforming density matrices on $\mathscr{H}_{in}$ into (normalized) density matrices on $\mathscr{H}_{out}$. Hence, our claim is that every instrument $\{\mathcal{P}_i\}_{i\in\mathsf{X}}$ can be decomposed as

$$\mathcal{P}_i = \mathcal{C}_i\mathcal{M}_i \qquad \forall i \in \mathsf{X}. \tag{3}$$

The proof is provided in the Supplemental Material. The decomposition of Eq. (3) is important because it states that any probabilistic process in quantum theory can be decomposed into a measurement, which modulates the amplitudes of the input state in a suitable basis, followed by a deterministic transformation depending on the outcome. In our case we consider measurements with two outcomes, referred to as *filters.* The filter induces a bifurcation of the wave-function and selects one particular branch, corresponding to one outcome with measurement operator M_{yes}.

Quantum metrology bounds. Here we sketch the proof that deterministic processes cannot reliably produce more than $O(N)$replicas (SQL), whereas probabilistic processes cannot produce more than $O(N^2)$ replicas (HL). We present the main ideas, while the full detail can be found in the supplementary material. Consider an exponential family of quantum states, of the form $\{|\psi_\theta\rangle = e^{i\theta H}|\psi\rangle\}$. When N identical copies are available, the quantum Cramér-Rao bound [7] states that the variance in the estimation of θ is lower bounded by $V_N(\theta) \geq \frac{4c}{N}$, for a constant c which can be set to $c = 1$ with a suitable change of parametrization. In this case, it is known that the bound can be achieved for large N. Suppose now that a *deterministic* process $\mathcal{C}$ produces $M = N + N^\alpha$, with an error vanishing in the limit of large N. To measure the error, we use the trace distance between the output of the process and the desired state of M perfect copies. For simplicity, we assume here that the error can be bounded as

$\|\mathcal{C}(|\psi_\theta\rangle\langle\psi_\theta|^{\otimes N}) - |\psi_\theta\rangle\langle\psi_\theta|^{\otimes M}\|_1 \leq M^{-\beta}$ for some $\beta \geq 1$ (this requirement will be lifted in the Supplemental Material, where we prove strong converses to the HL and SQL). We now use fact that the trace distance gives a bound on the total variation distance between the probability distributions arising from quantum measurements. This implies that the variance in the estimation of θ from the clones, denoted by $V_{out}(\theta)$, can be bounded as $V_{out}(\theta) \leq V_M(\theta) + \gamma M^{-\beta}$, for a suitable constant $\gamma > 0$. However, applying the quantum channel $\mathcal{C}$ cannot reduce the variance of the *optimal* estimation strategy, denoted by $V_N^{\min}(\theta)$. Therefore, we have the bound $V_N^{\min}(\theta) \leq V_{out}(\theta) \leq V_M(\theta) + \gamma M^{-\beta}$. Choosing the best estimation strategy, we then obtain

$$V_N^{\min}(\theta) \leq V_M^{\min}(\theta) + \gamma M^{-\beta} \tag{4}$$

For large N, we can replace the variances with the right hand side of the Cramér-Rao bound, thus obtaining

$$4/N \leq 4/(N+N^\alpha) + \gamma(N+N^\alpha)^{-\beta}. \tag{5}$$

Clearly, this implies $\alpha \leq 1$, because otherwise we would have a contradiction for large N. This establishes the standard quantum limit for reliable cloning. Moreover, we can also show that the reliability exponent β cannot be larger than 2. Indeed, by Taylor-expanding the r.h.s. of Eq. (5) we obtain the condition $\beta \leq 2-\alpha$, which implies $\beta \leq 2$. This means that no deterministic cloning process can produce copies with a trace distance vanishing faster than N^{-2}. Equivalently, the fidelity cannot approach 1 faster than N^{-4}. The difference with probabilistic processes is striking: remember that our filter allows to clone polarization states with an error that vanishes faster than any polynomial.

The derivation of the Heisenberg limit for reliable cloning follows the same lines sketched above. In the probabilistic case, the input state of the quantum channel $\mathcal{C}_{yes}$ is not the product state $|\psi_\theta\rangle^{\otimes N}$, but instead the entangled state $|\Phi_\theta^N\rangle = M_{yes}|\psi_\theta\rangle^{\otimes N}/\|M_{yes}|\psi_\theta\rangle^{\otimes N}\|$. Taking the average over all possible values of θ, the Heisenberg limit implies that the variance will be bounded as $V_{in,ave} \geq c/N^2$, for a suitable constant $c > 0$ (see the Supplemental Material for the proof). Choosing a cloning rate $\alpha > 1$ and a reliability exponent $\beta \geq 1$ and substituting the values in the quantum Cramér-Rao bound, Eq. (4) then becomes

$$cN^{-2} \leq 4N^{-\alpha} + O(N^{-\alpha}), \tag{6}$$

which implies $\alpha \leq 2$. Note that here there is no bound on the reliability rate β. This feature enables the exponential scaling of the fidelity demonstrated by our strategies.

Acknowledgments. This work is supported by the National Basic Research Program of China (973) 2011CBA00300 (2011CBA00301) and by the National Natural Science Foundation of China through Grants 61033001 and 61061130540 and by the 1000 Youth Fellowship Program of China. A particular acknowledgment goes to Xinhui Yang for drawing Figure 1.

[1] Wootters, W. & Zurek, W. A Single Quantum Cannot be Cloned. *Nature* **299,** 802-803(1982).

[2] Dieks, D. Communication by EPR devices. *Phys. Lett. A* **92,** 271-272(1982).

[3] Herbert, N. FLASH—A superluminal communicator based upon a new kind of quantum measurement. *Found. Phys.* **12,** 1171-1179(1982).

[4] This observation should be taken as a warning to remind that *stimulated emission is not a real physical process in the case of single photons.* Indeed, the unitarity of quantum mechanics implies that, by monitoring the environment, there is always the possibility to discover which probabilistic process took place on the system. Since for a single photon discovering if stimulated emission takes place is impossible in principle, stimulated emission cannot be a legitimate probabilistic process in this case.

[5] Wineland, D. J., Bollinger J. J., Itano, W. M. & Moore, F.L. Spin squeezing and reduced quantum noise in spectroscopy. *Phys. Rev. A* **46,** R6797-R6800(1992).

[6] Holland, M. J. & Burnett, K. Interferometric detection of optical phase shift at the Heisenberg limit. *Phys. Rev. Lett.* **71,** 1355(1993).

[7] Braunstein S. L. and Caves C. M., Statistical distance and the geometry of quantum states. *Phys. Rev. Lett.* **72,** 343943 (1994).

[8] Giovannetti, V., Lloyd, S. & Maccone, L. Quantum-enhanced measurements: beating the standard quantum limit. *Science* **306,** 1330-1336(2004).

[9] Giovannetti, V., Lloyd, S. & Maccone, L. Advances in quantum metrology. *Nature Photonics* **5,** 222-229(2011).

[10] Escher, B. M., de Matos Filho, R. L. & Davidovich, L. General framework for estimating the ultimate precision limit in noisy quantum-enhanced metrology. *Nature Physics* **7,** 406-411(2011).

[11] D'Ariano, G. & Macchiavello, C. Optimal phase-covariant cloning for qubits and qutrits. *Phys. Rev. A* **67,** 042306(2003).

[12] Stueckelberg, E. C. G. Quantum theory in real Hilbert space. *Helv. Phys. Acta* **33,** 458(1960).

[13] Hardy, L. & Wootters, W. Limited holism and real-vector-space quantum theory. *Found. Phys.* **3**, 454-473(2012).

[14] Wootters, W. Optimal Information Transfer and Real-Vector-Space Quantum Theory, arXiv:1301.2018.

[15] Xiang, G. Y., Ralph, T. C., Lund, A. P., Walk, N. & Pryde, G. J., Heralded noiseless linear amplification and distillation of entanglement, *Nature Photonics* **4**, 316319 (2010).

[16] Ferreyrol, F. et al. Implementation of a nondeterministic optical noiseless amplifier, *Phys. Rev. Lett.* **104**, 123603 (2010).

[17] Usuga, M. A. et al. Noise-powered probabilistic concen-

tration of phase information. *Nature Physics* **6**, 767771 (2010).

[18] Zavatta, A., Fiurášek, J. & Bellini, M. A high-fidelity noiseless amplifier for quantum light states. *N*ature Photonics **5**, 5260 (2011).

[19] Kocsis, S., Xiang, G. Y., Ralph, T. C. & Pryde, G. F. Heralded Noiseless Amplification of a Photon Polarization Qubit, *Nature Physics* **9**, 2328 (2013).

[20] Sherson, J. F., Krauter, H., Olsson, R. K., Julsgaard, B., Hammerer, K., Cirac, I. & Polzik, E. S., Quantum teleportation between light and matter, *Nature* **443**, 557-560 (2006).

[21] Duan, L. M. & Guo, G. C. A probabilistic cloning machine for replicating two non-orthogonal states, *Phys. Rev. Lett.* **80**, 49995002 (1998).

[22] J. Fiurášek, Phys. Rev. A **70**, 032308 (2004)

[23] Ralph, T. C. & Lund, A. P. Nondeterministic Noiseless Linear Amplification of Quantum Systems, *Quantum Communication Measurement and Computing Proceedings of 9th International Conference*, Ed. A. Lvovsky, 155-160 (AIP, New York 2009).

[24] Fiurasek, J., Optimal probabilistic estimation of quantum states. *New Jour. Phys.* **8,** 192(2006).

[25] Gendra, B., Ronco-Bonvehi, E., Calsamiglia, J., Muñoz-Tapia, R. & Bagan, E. Quantum metrology assisted with abstention, *Phys. Rev. Lett.* **110**, 100501 (2013).

Appendix A: Quantum Cramér-Rao bound for probabilistic metrology

Here we establish the Heisenberg limit for the probabilistic estimation of the parameter θ characterizing the state $|\psi_\theta\rangle = e^{i\theta H}|\psi\rangle$, showing that the variance can decrease at most as N^{-2} with the number N of input copies. In order to prove this result, we have first to establish a suitable Cramér-Rao bound for *probabilistic* quantum metrology. We will first by recalling some known facts about the Cramér-Rao bound in the usual deterministic scenario, and then use these facts to treat the probabilistic case.

In quantum metrology, the variance in estimation of parameter θ from a pure state $|\Psi_\theta\rangle$ is lower bounded by the Cramér-Rao bound

$$V(\theta) \geqslant \frac{1}{F_Q(|\Psi_\theta\rangle)} \qquad F_Q(|\Psi_\theta\rangle) := 4(\Delta H)^2, \tag{A1}$$

where $F_Q(|\Psi_\theta\rangle)$ called *quantum Fisher information*, H is the Hamiltonian that governs the evolution of $|\Psi_\theta\rangle$, namely $i\frac{d}{d\theta}|\Psi_\theta\rangle = H|\Psi_\theta\rangle$ and $(\Delta H)^2 := \langle\Psi_\theta|H^2|\Psi_\theta\rangle - \langle\Psi_\theta|H|\Psi_\theta\rangle^2$ is the variance of H.

For N copies of a pure state $|\psi_\theta\rangle = e^{i\theta H}|\psi\rangle$, the Hamiltonian is given by $H^{(N)} = \sum_{j=1}^N H_j$, where H_j denotes the Hamiltonian H acting on the Hilbert space of the j-th copy. Hence, the quantum Fisher information is given by $F_Q(|\psi_\theta\rangle^{\otimes N}) = 4N(\Delta H)^2$. Inserting this value in the Cramér-Rao bound one obtains the *standard quantum limit*, stating that the minimal variance for estimation using the product probe state $|\psi_\theta\rangle^{\otimes N}$ scales as $O(N^{-1})$.

We now consider the probabilistic case, where we want to estimate the parameter θ from the state $|\Psi_\theta\rangle$ using a filter. The joint state of the systems emerging from the filter is given by

$$|\Phi_\theta\rangle = \frac{M_{yes}|\Psi_\theta\rangle}{\|M_{yes}|\Psi_\theta\rangle\|}. \tag{A2}$$

To compute the Fisher information, we need to compute the variance of the Hamiltonian K_θ defined by $i\frac{d}{d\theta}|\Phi_\theta\rangle = K_\theta|\Phi_\theta\rangle$. Note that, in principle, K_θ can depend on θ, even if the original Hamiltonian H does not. By explicit calculation we obtain

$$K_\theta|\Phi_\theta\rangle = \left(M_{yes} H M_{yes}^{-1} + \frac{\langle\Psi_\theta|[H, M_{yes}^\dagger M_{yes}]|\Psi_\theta\rangle}{2\|M_{yes}|\Psi_\theta\rangle\|} \right)|\Phi_\theta\rangle. \tag{A3}$$

The quantum Fisher information at θ is then given by

$$F_Q(|\Phi_\theta\rangle) = \langle\Phi_\theta|K_\theta^\dagger K_\theta|\Phi_\theta^N\rangle - \left|\langle\Phi_\theta|K_\theta|\Phi_\theta\rangle\right|^2. \tag{A4}$$

Now, in order to minimize the variance one has to maximize the Fisher information over all possible filter operators M_{yes}. In this problem, there is an important caveat: as a matter of fact, for *every given value* of θ there one can choose a suitable M_{yes} that makes the quantum Fisher information as large as desired. This can be easily seen by choosing a parametrization where $\langle\Psi_\theta|H|\Psi_\theta\rangle = 0$ and defining $M_{yes,\theta} = \epsilon|\Psi_\theta\rangle\langle\Psi_\theta| - |\Psi_\theta^\perp\rangle\langle\Psi_\theta^\perp|$, where $|\Psi_\theta^\perp\rangle$ is the normalized vector proportional to $H|\Psi_\theta\rangle$. This choice gives a Fisher information scaling as $1/\epsilon^2$, which can be made arbitrarily large by choosing ϵ to be small. Hence, the usual Cramér-Rao bound (A1) would imply that probabilistic metrology can make the variance as small as desired. However, this is an artifact of the fact that the usual Cramér-Rao bound is a local bound: a probabilistic estimation scheme can have an unlimited precision *only at a particular value of* θ, which is the unknown parameter. Since θ is unknown, instead of minimizing the variance at a particular value of θ, it is more sensible to minimize the *average variance*.

$$V_{ave} := \int_\Theta \frac{d\theta}{2\pi}\; p(\theta|yes)\; V(\theta)$$

where Θ is the set of all possible values of θ, $p(\theta|yes)$ is the conditional probability distribution

$$p(\theta|yes) := p(\theta) \frac{\langle\Psi_\theta|M_{yes}^\dagger M_{yes}|\Psi_\theta\rangle}{\int_\Theta \frac{d\tau}{2\pi}\; p(\theta)\langle\Psi_\tau|M_{yes}^\dagger M_{yes}|\Psi_\tau\rangle},$$

and $p(\theta)$ is the prior distribution of θ. For simplicity, in the following we will consider the case of periodic evolutions, with $\Theta = [0, 2\pi)$, in which case we choose the prior to be uniform (i.e. $p(\theta) = 1$ for all θ). In the end of the section, we will discuss how to modify the argument in the non-periodic case $\Theta = \mathbb{R}$.

It is easy to see that the average variance is lower bounded by the inverse of the *averge quantum Fisher information* $F_{ave} := \int_\Theta d\theta/(2\pi)\; p(\theta|yes)\; F_Q(|\Psi_\theta\rangle)$. Indeed, using Eq. (A1) we have the

$$\begin{aligned} 1 &\leq \left[\int \frac{d\theta}{2\pi}\; p(\theta|yes)\; \sqrt{V(\theta) F_Q(|\Psi_\theta\rangle)}\right]^2 \\ &\leq \left[\int \frac{d\theta}{2\pi}\; p(\theta|yes)\; V(\theta)\right] \left[\int \frac{d\theta}{2\pi}\; p(\theta|yes)\; F_Q(|\Psi_\theta\rangle)\right], \end{aligned}$$

which implies the Cramér-Rao type bound $V_{ave} \geq \frac{1}{F_{Q,ave}}$. Minimizing the variance over all filter operators M_{yes} we then obtain

$$V_{ave}^{\min} \geq \frac{1}{F_{Q,ave}^{\max}} \tag{A5}$$

This form of the Cramér-Rao bound is the most suitable to study probabilistic metrology. We will now use it to show that for a family of states of the form (A2), with $|\Psi_\theta\rangle = |\psi_\theta\rangle^{\otimes N}$, $|\psi_\theta\rangle = e^{i\theta H}|\psi\rangle$, the average quantum Fisher information scales as $O(N^2)$. For simplicity, we will consider the case of phase estimation, where H has integer spectrum and θ is an angle $\theta \in [0, 2\pi)$, which is the case considered in the main text. Discarding the second term in the r.h.s. of Eq. (A4) and using Eq. (A3) we obtain the upper bound

$$\begin{aligned} F_Q(|\Phi_\theta\rangle) &\leqslant 4\langle\Phi_\theta|K_\theta^\dagger K_\theta|\Phi_\theta\rangle \\ &= 4\| K_\theta|\Phi_\theta\rangle\|^2 \\ &\leq 4\left(\left\| M_{yes} H^{(N)} M_{yes}^{-1}|\Phi_\theta\rangle\right\| + \left|\frac{\langle\Psi_\theta|[H^{(N)}, M_{yes}^\dagger M_{yes}]|\Psi_\theta\rangle}{2\|M_{yes}|\Psi_\theta\rangle\|^2}\right|\right)^2 \\ &\leqslant 4\left(\frac{\|M_{yes}H^{(N)}|\Psi_\theta\rangle\|}{\|M_{yes}|\Psi_\theta\rangle\|} + \frac{|\langle\Psi_\theta|M_{yes}^\dagger M_{yes} H^{(N)}|\Psi_\theta\rangle|}{\|M_{yes}|\Psi_\theta\rangle\|^2}\right)^2 \\ &\leqslant 4\left(\frac{2\|M_{yes}H^{(N)}|\Psi_\theta\rangle\|}{\|M_{yes}|\Psi_\theta\rangle\|}\right)^2 \\ &= \frac{16\;\langle\Psi_\theta|\; H^{(N)}\; M_{yes}^\dagger M_{yes}\; H^{(N)}\;|\Psi_\theta\rangle}{\langle\Psi_\theta|\; M_{yes}^\dagger M_{yes}|\Psi_\theta\rangle}. \end{aligned}$$

Taking the average over θ we then have

$$
\begin{aligned}
F^{\max}_{Q,ave} &\le \max_{M_{yes}} \int_0^{2\pi} d\theta\ p(\theta|yes) \frac{16\ \langle\Psi_\theta|\ H^{(N)}\ M^\dagger_{yes}M_{yes}\ H^{(N)}\ |\Psi_\theta\rangle}{\langle\Psi_\theta|\ M^\dagger_{yes}M_{yes}|\Psi_\theta\rangle} \\
&= \max_{M_{yes}} \frac{16\ \int \frac{\mathrm{d}\theta}{2\pi}\ p(\theta)\ \langle\Psi_\theta|\ H^{(N)}\ M^\dagger_{yes}M_{yes}\ H^{(N)}\ |\Psi_\theta\rangle}{\int \frac{\mathrm{d}\tau}{2\pi}\ p(\tau)\ \langle\Psi_\tau|\ M^\dagger_{yes}M_{yes}\ |\Psi_\tau\rangle} \\
&= \max_{M_{yes}} \frac{16\ \langle\Psi|\ H^{(N)}\ \left(\int \frac{\mathrm{d}\theta}{2\pi}\ p(\theta)\ e^{-i\theta H^{(N)}} M^\dagger_{yes}M_{yes}\ e^{i\theta H^{(N)}}\right)\ H^{(N)}\ |\Psi\rangle}{\langle\Psi|\ \left(\int \frac{\mathrm{d}\tau}{2\pi}\ p(\tau)\ e^{-i\tau H^{(N)}} M^\dagger_{yes}M_{yes} e^{i\tau H^{(N)}}\right) |\Psi\rangle} \\
&= \max_{\widetilde{M}_{yes}} \frac{16\ \langle\Psi|\ H^{(N)}\ \widetilde{M}^\dagger_{yes}\widetilde{M}_{yes}\ H^{(N)}\ |\Psi\rangle}{\langle\Psi|\ \widetilde{M}^\dagger_{yes}\widetilde{M}_{yes}|\Psi\rangle} \qquad \widetilde{M}_{yes} := \sqrt{\int \frac{\mathrm{d}\theta}{2\pi}\ p(\theta)\ e^{-i\theta H^{(N)}} M^\dagger_{yes}M_{yes}\ e^{i\theta H^{(N)}}}
\end{aligned}
$$

Noting that $\widetilde{M}_{yes}$ commutes with $H^{(N)}$, we then obtain

$$
\begin{aligned}
F^{\max}_{Q,ave} &\le \max_{M_{yes}:\ [M_{yes}, H^{(N)}]=0} \frac{16\ \langle\Psi|\ H^{(N)}\ M^\dagger_{yes}M_{yes}\ H^{(N)}\ |\Psi\rangle}{\langle\Psi|\ M^\dagger_{yes}M_{yes}|\Psi\rangle} \\
&= \max_{M_{yes}:\ [M_{yes}, H^{(N)}]=0} \frac{16\ \langle\Psi|\ M^\dagger_{yes}H^{(N)}H^{(N)}\ M_{yes}\ |\Psi\rangle}{\langle\Psi|\ M^\dagger_{yes}M_{yes}|\Psi\rangle} \\
&\le 16\|H^{(N)}\|^2_\infty,
\end{aligned}
$$

where $\|H^{(N)}\|_\infty$ denotes the operator norm of $H^{(N)}$, equal to the maximum of the modulus of its eigenvalues. Since $H^{(N)} = \sum_{j=1}^N H_j$ we have that $\|H^{(N)}\|_\infty$ scales linearly with N, thus giving the bound $F^{\max}_{Q,ave} \le O(N^2)$, and therefore $V^{\min}_{ave} \ge O(N^{-2})$.

In the non-periodic case $\theta \in \mathbb{R}$, all the arguments can be repeated by choosing a Gaussian prior $p_\lambda(\theta) = \sqrt{2\pi/\sigma^2}e^{-\theta^2/(2\sigma^2)}$, the only difference being that now the operator $\widetilde{M}_{yes}$ does not commute with $H^{(N)}$. However, we can always replace $\widetilde{M}_{yes}$ by its decohered version, where all the off-diagonal elements in the eigenbasis of $H^{(N)}$ are replaced by zero. Since these off-diagonal elements are exponentially small in the variance σ^2, the corresponding correction can be made arbitrarily small, leading to the bound $F^{\max}_{Q,ave} \le 16\|H^{(N)}\|^2_\infty + \epsilon$ with arbitrary small ϵ.

Appendix B: Decomposition of quantum instruments

Here we prove that any quantum instrument $\{\mathcal{P}_i\}_{i\in\mathsf{X}}$ can be decomposed into a pure measurement on the input system, followed by a deterministic process conditional to the outcome of the measurement.

The proof is simple: let $\mathcal{P}^\dagger_i$ be the adjoint of the map $\mathcal{P}_i$, defined by $\mathrm{Tr}[\mathcal{P}^\dagger_i(A)\rho] = \mathrm{Tr}[A\mathcal{P}_i(\rho)]$ for every operator A and for every density matrix ρ. Define the operator $M_i := \sqrt{\mathcal{P}^\dagger_i(I)}$ and the map $\mathcal{L}_i$ by the relation $\mathcal{L}_i(\rho) := \mathcal{P}_i(M_i^{-1}\rho M_i^{-1})$, where M_i^{-1} is the inverse of M_i on its support. If M_i is invertible, then $\mathcal{C}_i := \mathcal{L}_i$ will be trace-preserving (and hence a quantum channel). In this case, by definition we will have $\mathcal{P}_i = \mathcal{C}_i\mathcal{M}_i$. If M_i is not invertible, we can define the trace-preserving map $\mathcal{C}_i(\rho) := \mathcal{L}_i(\rho) + Q_0\rho Q_0$, where Q_0 is the projector on the kernel of M_i. Again, the definition implies $\mathcal{P}_i = \mathcal{C}_i\mathcal{M}_i$. Hence, we established the decomposition $\mathcal{P}_i = \mathcal{C}_i\mathcal{M}_i, \quad \forall i \in \mathsf{X}$.

Appendix C: General facts about quantum cloning

Here we give the general formulation of the problem of optimal cloning in the presence of symmetry, taking into account arbitrary probabilistic processes. The general facts introduced here will be specialized to the case of cloning of states of the form $|\psi_\theta\rangle = e^{i\theta H}|\psi\rangle$ in the next section.

1. Probabilistic cloning in the presence of symmetries

We consider probabilistic cloning processes designed to produce approximate copies of a pure state of the form $|\psi_g\rangle := U_g|\psi\rangle$, where $|\psi\rangle \in \mathscr{H}$ is a fixed unit vector and $\{U_g\}$ is a unitary representation of a group G on the

Hilbert space of the system. A probabilistic cloning process is described by a quantum instrument $\{\mathcal{P}_{yes}, \mathcal{P}_{no}\}$ consisting of two quantum operations (completely positive trace non-increasing maps) corresponding to the successful and unsuccessful instances of the process, respectively. For a cloning machine designed to transform N input copies into M approximate copies, the probability of success is given by $p(yes|g) := \mathrm{Tr}\{\mathcal{P}_{yes}[(|\psi_g\rangle\langle\psi_g|)^{\otimes N}]\}$ and the state of the M output systems is $\rho'_g := \mathcal{P}_{yes}[(|\psi_g\rangle\langle\psi_g|)^{\otimes N}]/\,\mathrm{Tr}\{\mathcal{P}_{yes}[(|\psi_g\rangle\langle\psi_g|)^{\otimes N}]\}$.

When the group G is compact, it is natural to assume that the prior probability of the states $|\psi_g\rangle$ is uniform and given by the haar measure $\underset{\cdot}{\mathrm{g}}$. In this case, the average fidelity of the output of the device with the desired state $|\psi_g\rangle^{\otimes M}$, conditionally to the successful realization of the process, is given by

$$F = \int \underset{\cdot}{\mathrm{g}}\ p(g|yes)\ \langle\psi_g|^{\otimes M}\rho'_g|\psi_g\rangle^{\otimes M} = \frac{\int \underset{\cdot}{\mathrm{g}}\ \langle\psi_g|^{\otimes M}\mathcal{P}_{yes}[(|\psi_g\rangle\langle\psi_g|)^{\otimes N}]|\psi_g\rangle^{\otimes M}}{\int \underset{\cdot}{\mathrm{g}}\ \ \mathrm{Tr}\{\mathcal{P}_{yes}[(|\psi_g\rangle\langle\psi_g|)^{\otimes N}]\}}. \tag{C1}$$

From the form of Eq. (C1) is is clear that the search for the optimal quantum operation $\mathcal{P}_{yes}$ can be restricted without loss of generality to the set of maps satisfying the covariance condition $\mathcal{P}_{yes}\mathcal{U}_g^{\otimes N} = \mathcal{U}_g^{\otimes M}\mathcal{P}_{yes}, \forall g \in G$, where $\mathcal{U}_g$ is the unitary channel defined by $\mathcal{U}_g(\rho) = U_g\rho U_g^\dagger$. When the group G is non-compact, restricting to covariant processes means requiring that both the probability of success and the fidelity are independent of the unknown group element g.

Decomposing the covariant quantum operation $\mathcal{P}_{yes}$ as $\mathcal{P}_{yes} = \mathcal{C}_{yes}\mathcal{M}_{yes}$ where $\mathcal{M}_{yes}(\rho) = M_{yes}\rho M_{yes}^\dagger$ is a pure measurement and $\mathcal{C}_{yes}$ is a quantum channel (completely positive trace-preserving map) we have that the latter must satisfy the symmetry conditions

$$[M_{yes}, U_g^{\otimes N}] = 0 \qquad\qquad \forall g \in G \tag{C2}$$

$$\mathcal{C}_{yes}\mathcal{U}_g^{\otimes N} = \mathcal{U}_g^{\otimes M}\mathcal{C}_{yes} \qquad\qquad \forall g \in G. \tag{C3}$$

Using the decomposition $\mathcal{P}_{yes} = \mathcal{C}_{yes}\mathcal{M}_{yes}$ and the above symmetry conditions, one can transform Eq. (C1) into the form

$$F = \langle\psi|^{\otimes M}\mathcal{C}_{yes}(|\Psi^N\rangle\langle\Psi^N|)|\psi\rangle^{\otimes M}, \tag{C4}$$

where $|\Psi^N\rangle$ is the conditional state

$$|\Psi^N\rangle := \frac{M_{yes}|\psi\rangle^{\otimes N}}{\|M_{yes}|\psi\rangle^{\otimes N}\|}. \tag{C5}$$

Note also that the symmetry condition of Eq. (C2) implies that the probability that the input system pass the filter is independent of g and is given by $p_{yes} = \|M_{yes}|\psi\rangle^{\otimes N}\|^2$.

When $\{U_g,\ g \in \mathsf{G}\}$ are *all* the unitary operators on the Hilbert space of the system, any operator M_{yes} that commutes with all unitaries will act on $|\psi_g\rangle^{\otimes N}$ as a multiple of the identity, due to the Schur's lemma. Hence, the action of the filter will not change the input state: $|\Phi_g^N\rangle = M_{yes}|\psi_g\rangle^{\otimes N}/\||\psi_g\rangle^{\otimes N}\| = |\psi_g\rangle^{\otimes N}$ for every value of g. We conclude that in this case the symmetry of the set of states inhibites super-replication, by enforcing the standard quantum limit. The same conclusion is reached if $\{U_g, g \in \mathsf{G}\}$ is an irreducible representation and if $|\psi\rangle$ is a lowest weight-vector, because also in this case the operator M_{yes} acts as a multiple of the identity on the states $|\psi_g\rangle^{\otimes N}$, inhibiting super-replication. For example, super-replication is impossible for the arbitrary coherent states $|\alpha\rangle = e^{-|\alpha|^2/2}\sum_{n=0}^\infty \alpha^n/\sqrt{n!}|n\rangle$ of the electromagnetic field (as opposed to the case of coherent states with fixed amplitude).

Appendix D: Super-replication of Hamiltonian-generated quantum states

Consider the case the group is either $U(1) \simeq [0, 2\pi)$, in the case of periodic evolutions, or $\mathbb{R}$, in the case of aperiodic evoultions. The representation of interest is $U_\theta = e^{i\theta H}$, and the input state is $|\psi\rangle = \sum_{E\in\mathsf{Spec}(H)}\sqrt{p_E}|E\rangle$, where $\mathsf{Spec}(H)$ denotes the spectrum of H and $|E\rangle \in \mathscr{H}$ is a normalized eigenvector corresponding to the eigenvalue E. We will assume without loss of generality that the vectors $\{|E\rangle \mid E \in \mathsf{Spec}(H)\}$ span $\mathscr{H}$ and that $\langle\psi|H|\psi\rangle = 0$. When N copies are given, we write $|\psi\rangle^{\otimes N} = \sum_{E\in\mathsf{Spec}(H^{(N)})}\sqrt{p_{N,E}}|N, E\rangle$, where $|N, E\rangle \in \mathscr{H}^{\otimes N}$ is a normalized eigenvector of $H^{(N)}$ corresponding to the eigenvalue E. By the central limit theorem, for large N the probability distribution $p_{N,E}$ converges in expectation values to the Gaussian $g_{N,E} = e^{-E^2/(2N\langle H^2\rangle)}/\sqrt{2\pi N\langle H^2\rangle}$.

1. Strong converse of the quantum metrology limits

Here we consider super-replication processes producing $M = cN^{1+\epsilon}$, $\epsilon > 0$, $c > 0$ output copies. We determine the ultimate limits to super-replication, showing that

1. Every super-replication process with success probability $p_{yes} \geq ae^{-bN^x}$ for some $a, b > 0$ and $x < \epsilon$ must have vanishing fidelity. In particular, this implies that every deterministic process producing more than $O(N)$ output copies must have vanishing fidelity.
2. Every probabilistic process producing more than cN^2 output copies for some $c > 0$ must have vanishing fidelity.

The above results provide a strong converse of the standard quantum limit and of the Heisenberg limit, respectively.

Let us introduce the Choi operator of the channel $\mathcal{C}_{yes}$, denoted by C_{yes}. In terms of the Choi operator, the symmetry condition of Eq. (C3) becomes $[C_{yes}, U_\theta^{\otimes M} \otimes U_{-\theta}^{\otimes N}] = 0$, $\forall \theta \in [0, 2\pi)$. Hence, we have that C_{yes} can be written in the block diagonal form $C_{yes} = \bigoplus_{\mu \in \mathsf{Spec}\left(H^{(M)} - H^{(N)}\right)} C_\mu$, with each block in the form of

$$C_\mu = \sideset{}{'}\sum_{E, E' \in \mathsf{Spec}\left(H^{(N)}\right)} [C_\mu]_{E,E'} \ \ |M, E' + \mu\rangle\langle M, E + \mu| \otimes |N, E\rangle\langle N, E'| \,, \tag{D1}$$

where the summation $\sum'$ is restricted to the values of E (E') such that $E + \mu$ $(E' + \mu)$ belongs to $\mathsf{Spec}(H^{(M)})$.

The trace-preserving condition on $\mathcal{C}_{yes}$ can be re-written as

$$\sideset{}{'}\sum_{\mu \in \mathsf{Spec}\left(H^{(M)} - H^{(N)}\right)} [C_\mu]_{E,E} = 1 \qquad \forall E \in \mathsf{Spec}\left(H^{(N)}\right), \tag{D2}$$

where the summation is restricted to the values of μ such that $E + \mu$ belongs to $\mathsf{Spec}(H)$. The fidelity in Eq. (C4) becomes

$$F = \langle\psi|^{\otimes M} \langle\Psi^N| C_{yes} |\psi\rangle^{\otimes M} |\Psi^N\rangle,$$

to be maximized over the set of positive operators C_{yes} satisfying the constraints of Eqs. (D1) and (D2). Expanding the states $|\psi\rangle^{\otimes M}$ and $|\Psi^N\rangle$ as

$$|\psi\rangle^{\otimes M} = \sum_{E \in \mathsf{Spec}\left(H^{(M)}\right)} \sqrt{p_{M,E}} |M, E\rangle \tag{D3}$$

$$|\Psi^N\rangle = \sum_{E \in \mathsf{Spec}\left(H^{(N)}\right)} \Psi_E^N |N, E\rangle, \tag{D4}$$

(for some suitable coefficients $\{\Psi_E^N\}$ satisfying $\sum_{E \in \mathsf{Spec}\left(H^{(N)}\right)} |\Psi_E^N|^2 = 1$), we then obtain

$$F = \sum_\mu \left(\sideset{}{'}\sum_{E,E'} [C_\mu]_{E,E'} \sqrt{p_{M,E+\mu}} \Psi_E^{N*} \sqrt{p_{M,E'+\mu}} \Psi_{E'}^N \right). \tag{D5}$$

The positivity of the operator C_{yes} (and hence of the matrices C_μ) implies the inequality $|[C_\mu]_{E,E'}| \leqslant \sqrt{c_E^\mu c_{E'}^\mu}$, $c_E^\mu := [C_\mu]_{E,E}$, which leads to the bound

$$F \leq \sum_\mu \left[\sideset{}{'}\sum_E \sqrt{c_E^\mu \ p_{M,E+\mu}} |\Psi_E^N| \right]^2 . \tag{D6}$$

Picking $0 < \delta < \epsilon$ and defining $E_\delta = 2\sqrt{N^{1+\delta} \langle H^2 \rangle}$ we get

$$\begin{aligned} F &\leq \sum_\mu \left(\sideset{}{'}\sum_{|E| \leq E_\delta} \pi_E \sqrt{\frac{c_E^\mu p_{M,E+\mu} p_{N,E}}{p_{yes}}} + \sideset{}{'}\sum_{|E| > E_\delta} \pi_E \sqrt{\frac{c_E^\mu p_{M,E+\mu} p_{N,E}}{p_{yes}}} \right)^2 \\ &\leq \left[\sqrt{\sum_\mu \left(\sideset{}{'}\sum_{|E| \leq E_\delta} \pi_E \sqrt{\frac{c_E^\mu p_{M,E+\mu} p_{N,E}}{p_{yes}}} \right)^2} + \sqrt{\sum_\mu \left(\sideset{}{'}\sum_{E > E_\delta} \pi_E \sqrt{\frac{c_E^\mu p_{M,E+\mu} p_{N,E}}{p_{yes}}} \right)^2} \right]^2 . \end{aligned} \tag{D7}$$

For large N, the first term in Eq. (D7) is vanishes whenever $\delta < \epsilon$: indeed, one has

$$\begin{aligned}
\lim_{N\to\infty}\sum_{\mu}\left(\sideset{}{'}\sum_{|E|\le E_\delta}\pi_E\sqrt{\frac{c_E^{\mu}p_{M,E+\mu}p_{N,E}}{p_{yes}}}\right)^2 &\le \lim_{N\to\infty}\sum_{\mu}\left(\sideset{}{'}\sum_{|E|\le E_\delta}p_{M,E+\mu}\right)\left(\sideset{}{'}\sum_{|E|\le E_\delta}\frac{\pi_E^2 c_E^{\mu}p_{N,E}}{p_{yes}}\right)\\
&\le \lim_{N\to\infty}\sum_{\mu}\left(\sideset{}{'}\sum_{|E|\le E_\delta}p_{M,E}\right)\left(\sideset{}{'}\sum_{|E|\le E_\delta}\frac{\pi_E^2 c_E^{\mu}p_{N,E}}{p_{yes}}\right)\\
&\le \lim_{N\to\infty}\left(\sum_{|E|\le E_\delta}p_{M,E}\right) = 0,
\end{aligned}$$

where the last equality is due to the fact that $p_{M,E}$ converges to a Gaussian with variance of order $\sqrt{N^{\epsilon}}$, and the sum is over a set of negligible size $\sqrt{N^{\delta}}$.

If $\epsilon > 1$, this is enough to prove that the fidelity vanishes, no matter how small the probability is: indeed, choosing $1 < \delta < \epsilon$, we have that $|E| \le E_\delta$ for every $E \in \mathsf{Spec}(H^{(N)})$ and therefore, we do not have the second term in Eq. (D7). Hence, we have proved a strong converse of the Heisenberg limit: every process producing more than $O(N^2)$ output copies must have vanishing fidelity.

If $\epsilon < 1$, then the second term in Eq. (D7) can be upper bounded as

$$\begin{aligned}
\lim_{N\to\infty}\sum_{\mu}\left(\sideset{}{'}\sum_{|E|> E_\delta}\pi_E\sqrt{c_E^{\mu}p_{M,E+\mu}p_{N,E}}\right)^2 &\le \lim_{N\to\infty}\frac{g_{N,E_\delta}}{p_{yes}}\sum_{\mu}\left(\sideset{}{'}\sum_{|E|> E_\delta}\sqrt{\frac{c_E^{\mu}p_{M,E+\mu}}{p_{yes}}}\right)^2\\
&\le \lim_{N\to\infty}\frac{g_{N,E_\delta}}{p_{yes}}\sum_{\mu}\left(\sideset{}{'}\sum_{E> E_\delta}c_E^{\mu}\right)\left(\sideset{}{'}\sum_{E> E_\delta}p_{M,E+\mu}\right)\\
&\le \lim_{N\to\infty}\frac{g_{N,E_\delta}}{p_{yes}}\left(\sideset{}{'}\sum_{E> E_\delta}1\right)\\
&\le \lim_{N\to\infty}\frac{g_{N,E_\delta}}{p_{yes}}\binom{N+|\mathsf{Spec}(H)|-1}{|\mathsf{Spec}(H)|-1}\\
&\le \frac{1}{\sqrt{2\pi\langle H^2\rangle}}\lim_{N\to\infty}\frac{e^{-N^{\delta}}}{p_{yes}}.
\end{aligned}$$

Now, if $p_{yes} \ge a e^{-bN^{x}}$ for some $a, b > 0$ and $x < \epsilon$, then choosing δ such that $x < \delta < \epsilon$ we have $\lim_{N\to\infty}\frac{e^{-N^{\delta}}}{p_{yes}} = 0$, implying that the fidelity vanishes. This proves a strong converse of the standard quantum limit: every deterministic process producing more than $O(N)$ output copies must have vanishing fidelity.

2. Performances of the filter

Here we evaluate the fidelity and the probability of success of the super-replication strategy proposed in the main text. Let us first recall the ingredients: The probabilistic filter is given by

$$M_{yes} = \sum_{E\in\mathsf{Spec}(H^{(N)})}\pi_E|N,E\rangle\langle N,E| \qquad \pi_E = \gamma\sqrt{\frac{p_{M,E+E_0}}{p_{N,E}}}, \tag{D8}$$

where E_0 be the smallest energy shift such that $E + E_0$ belongs to $\mathsf{Spec}(H^{(M)})$ for every $E \in \mathsf{Spec}(H^{(N)})$ and γ is a suitable constant. The deterministic channel $\mathcal{C}_{yes}$ is given by the isometry

$$V = \sum_{E\in\mathsf{Spec}(H^{(N)})}|M,E+E_0\rangle\langle N,E|. \tag{D9}$$

The fidelity is then given by

$$F = \sum_{E\in\mathsf{Spec}(H^{(N)})}p_{M,E+E_0} \ge \sum_{|E|\le NE_{\min}}p_{M,E+E_0}, \qquad E_{\min} := \min_{E\in\mathsf{Spec}(H),E\neq 0}|E|. \tag{D10}$$

In the limit for large M, the shift E_0 tends to zero and, by the central limit theorem, the r.h.s. converges to $\mathrm{erf}(NE_{\min}/\sqrt{2M\langle H^2\rangle})$, where erf denotes the error function. Lower-bounding the error function we then get

$$F \geq 1 - \sqrt{\frac{2M\langle H^2\rangle}{\pi E_{\min}^2 N^2}}\exp\left(-\frac{N^2E_{\min}^2}{2M\langle H^2\rangle}\right) \tag{D11}$$

Let us now evaluate the probability of success, given by $p_{yes} = \gamma^2\sum_{E\in\mathsf{Spec}(H^{(N)})} p_{M,E} \leq \gamma^2$. Since π_E in Eq. (D8) must be smaller than 1 for every E, the largest possible value for the constant γ is $\gamma = \min_{E\in\mathsf{Spec}(H^{(N)})}\sqrt{\frac{p_{N,E}}{p_{M,E+E_0}}}$. In the limit of large N, this gives $p_{yes} \leq \exp\left(-\frac{N\|H\|_\infty^2}{4\langle H^2\rangle}\right)$

3. Increasing the probability of success

Here we devise super-replication processes that produce $M = cN^{1+\epsilon}$ copies for every desired $\epsilon \in [0,1)$ and $c > 0$, with fidelity approaching 1 in the limit $N \to \infty$, and with success probability lower bounded as $p_{yes} \geq e^{-N^\epsilon f(N)}$, where $f(N)$ is any fixed positive function such that $\lim_{N\to\infty} f(N) = \infty$. For $\epsilon > 0$, this proves in particular that one can achieve success probability larger than e^{-N^δ} for every $\delta > \epsilon$. For $\epsilon = 0$, it proves in particular that one can achieve success probability larger than $1/N^\delta$, for every $\delta > 0$.

Consider the filter given by

$$M_{yes} = \sum_{E\in\mathsf{Spec}(H^{(N)})} \pi_E |N,E\rangle\langle N,E|, \qquad \pi_E = \begin{cases} \gamma\sqrt{\frac{p_{M,E+E_0}}{p_{N,E}}} & |E| \leq E_{f(N)} := \sqrt{2M\langle H^2\rangle f(N)} \\ \gamma := \min_{|E|\leq E_{f(N)}}\sqrt{\frac{p_{N,E}}{p_{M,E+E_0}}} & \\ 1 & |E| > E_{f(N)}. \end{cases} \tag{D12}$$

The combination of the filter M_{yes} with the isometry V of Eq. (D9) gives a probabilistic process with fidelity lower bounded by $F \geq \sum_{|E|\leq E_{f(N)}} p_{M,E+E_0}$, which in the limit of large N, the r.h.s. converges to the error function $\mathrm{erf}\left(\sqrt{f(N)}\right)$, giving the bound

$$F \geq 1 - \frac{1}{\sqrt{\pi f(N)}}\exp[-f(N)]. \tag{D13}$$

The probability of success of the above cloning process is lower bounded by

$$p_{yes} \geq \sum_{|E|\leq E_{f(N)}} \pi_E^2\, p_{N,E} = \gamma^2 \sum_{|E|\leq E_{f(N)}} p_{M,E+E_0}. \tag{D14}$$

In the limit $N \to \infty$, the r.h.s. converges to $g_{N,E_{f(N)}}/g_{M,E_{f(N)}}\,\mathrm{erf}\left(\sqrt{f(N)}\right) \geq e^{-N^\epsilon f(N)}$.